



Cambridge International AS & A Level

CANDIDATE
NAMECENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

CHEMISTRY

9701/42

Paper 4 A Level Structured Questions

May/June 2026

2 hours

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen. Do **not** use correction fluid or tape.
- Do **not** write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 100.
- The number of marks for each question or part question is shown in brackets [].
- The Periodic Table is printed in the question paper.
- Important values, constants and standards are printed in the question paper.

This document has **28** pages.

- 1 (a) (i) Transition elements have variable oxidation states and form complex ions.

State **two** other typical chemical properties of a transition element.

- 1
- 2 [1]

- (ii) Explain why transition elements have variable oxidation states.

.....

.....

..... [1]

- (b) An aqueous solution of chromium(III) ions, Cr^{3+} , contains the $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$ complex ion.

- (i) Complete Figure 1.1 to show the electronic configuration for Cr^{3+} .




Figure 1.1

[1]

- (ii) State the type of bonding that exists between H_2O and Cr^{3+} in $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$.

..... [1]

- (c) Explain why transition elements form complex ions.

.....

.....

..... [1]

- (d) $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$ can be converted into the $[\text{Cr}(\text{H}_2\text{O})_3(\text{OH})_3]$ complex.

- (i) Suggest a suitable reagent for this conversion. State the type of reaction.

reagent

type of reaction [1]



- (ii) $[\text{Cr}(\text{H}_2\text{O})_3(\text{OH})_3]$ reacts with an excess of $\text{NH}_3(\text{aq})$ to form $[\text{Cr}(\text{NH}_3)_6]^{3+}$.

Complete the equation for this reaction. State the type of reaction.



type of reaction $\dots\dots\dots$

[2]

- (e) $[\text{Cr}(\text{H}_2\text{O})_3(\text{OH})_3]$ shows stereoisomerism.

Draw three-dimensional diagrams in Figure 1.2 to suggest the **two** stereoisomers of $[\text{Cr}(\text{H}_2\text{O})_3(\text{OH})_3]$.




Figure 1.2

[2]

- (f) (i) An aqueous solution of iron(III) ions, Fe^{3+} , contains the $[\text{Fe}(\text{H}_2\text{O})_6]^{3+}$ complex ion. This solution is acidic in a similar way to an aqueous solution of Al^{3+} .

Suggest why a solution of Fe^{3+} is acidic. Include an equation in your answer.

.....
.....
..... [2]

- (ii) An aqueous solution of iron(II) ions, Fe^{2+} , contains the $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$ complex ion. This solution is also acidic.

Predict how the pH of $1.00 \text{ mol dm}^{-3} \text{ Fe}^{2+}(\text{aq})$ differs from the pH of $1.00 \text{ mol dm}^{-3} \text{ Fe}^{3+}(\text{aq})$. Explain your answer.

.....
.....
..... [1]

- (g) Suggest why solutions of silver(I) salts are colourless.

.....
.....
.....
..... [2]




- (h) A mixture contains sodium ethanedioate, $\text{Na}_2\text{C}_2\text{O}_4$, ethanedioic acid dihydrate, $\text{H}_2\text{C}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$, and an unreactive solid.


5.00 g of this mixture is dissolved in distilled water and made up to 500 cm^3 in a volumetric flask to form solution **A**.

Two different titration experiments are carried out using **A**.

In the first experiment, 25.0 cm^3 of **A** requires 21.80 cm^3 of acidified $0.0250\text{ mol dm}^{-3}$ MnO_4^- (aq) to reach the end-point.



In the second experiment, 25.0 cm^3 of **A** requires 13.60 cm^3 of 0.150 mol dm^{-3} OH^- (aq) to reach the end-point.



Calculate the percentage by mass of $\text{Na}_2\text{C}_2\text{O}_4$ in the mixture.

$[M_r: \text{Na}_2\text{C}_2\text{O}_4, 134.0]$

percentage by mass of $\text{Na}_2\text{C}_2\text{O}_4 = \dots\dots\dots\%$ [5]

[Total: 20]

- 2 Group 2 carbonates decompose on heating to form the corresponding metal oxide and carbon dioxide.

- (a) The mechanism for this decomposition involves the carbonate ion breaking down to form carbon dioxide and oxide ions.

Add **two** curly arrows to the carbonate ion in Figure 2.1 to suggest a mechanism for this decomposition.




Figure 2.1

[1]

- (b) The radii of some cations are shown in Table 2.1.

Table 2.1

cation	Ca ²⁺	Cu ²⁺	Pb ²⁺
radius/nm	0.099	0.073	0.120

Use the data in Table 2.1 to predict the trend in thermal stability of the metal carbonates containing these cations.

Explain your answer.

..... > >
 most stable least stable

explanation

.....

.....

.....

.....

[2]



(c) Suggest why the solubility of the Group 2 sulfates decreases down the group.

.....

.....

.....

.....

.....

..... [3]

(d) Magnesium hydroxide, $\text{Mg}(\text{OH})_2$, is sparingly soluble in water at 298 K.

(i) Write the expression for the solubility product, K_{sp} , of $\text{Mg}(\text{OH})_2$. State the units for K_{sp} .

$$K_{\text{sp}} =$$

units [2]

(ii) A saturated solution of $\text{Mg}(\text{OH})_2$ has a pH of 10.35 at 298 K.

Calculate the concentration of hydroxide ions, OH^- , in mol dm^{-3} , in this solution.


concentration of OH^- = mol dm^{-3} [1]

(iii) Use your answers to **2(d)(i)** and **2(d)(ii)** to calculate the value of K_{sp} for $\text{Mg}(\text{OH})_2$.

K_{sp} for $\text{Mg}(\text{OH})_2$ = [1]

[Total: 10]

- 3 (a) Ethanal, CH_3CHO , reacts with methanol, CH_3OH , under acidic conditions to form compound **B**, as shown in reaction 1.



The initial rate of reaction is investigated using different starting concentrations of CH_3CHO , CH_3OH and H^+ . The results obtained are shown in Table 3.1.

Table 3.1

experiment	$[\text{CH}_3\text{CHO}]$ $/\text{mol dm}^{-3}$	$[\text{CH}_3\text{OH}]$ $/\text{mol dm}^{-3}$	$[\text{H}^+]$ $/\text{mol dm}^{-3}$	initial rate $/\text{mol dm}^{-3} \text{s}^{-1}$
1	0.500	0.040	0.100	2.500×10^{-3}
2	0.500	0.060	0.100	3.750×10^{-3}
3	0.250	0.120	0.300	1.125×10^{-2}
4	0.250	0.120	0.400	1.500×10^{-2}

- (i) Use the information in Table 3.1 to deduce the rate equation for this reaction.

Explain your reasoning. State clearly which experiments you use in your explanation.

.....

.....

.....

.....

.....

.....

.....

.....

rate =

[4]

- (ii) State the units of the rate constant, k , for this reaction.

..... [1]



(iii) Another experiment to investigate reaction 1 is carried out as shown in Table 3.2.


Table 3.2

experiment	$[\text{CH}_3\text{CHO}]/\text{mol dm}^{-3}$	$[\text{CH}_3\text{OH}]/\text{mol dm}^{-3}$	$[\text{H}^+]/\text{mol dm}^{-3}$
5	0.250	0.250	0.250

Calculate the initial rate of experiment 5.

initial rate = $\text{mol dm}^{-3} \text{s}^{-1}$ [1]

- (b) The reaction of propanone, CH_3COCH_3 , with bromine, Br_2 , in the presence of an acid catalyst is shown in reaction 2.



The rate of reaction is first order with respect to $[\text{CH}_3\text{COCH}_3]$ and first order with respect to $[\text{H}^+]$ under certain conditions.

A three-step mechanism for reaction 2 is suggested.

Complete Table 3.3 to suggest steps 1 and 3 of the mechanism.

Table 3.3

step 1 (fast)	
step 2 (slow)	$\text{H}_3\text{C}-\overset{\overset{\text{OH}^+}{\parallel}}{\text{C}}-\text{CH}_3 \longrightarrow \text{H}_3\text{C}-\overset{\overset{\text{OH}}{\mid}}{\text{C}}=\text{CH}_2 + \text{H}^+$
step 3 (fast)	

[2]



(c) Catalysts can be described as homogeneous or heterogeneous.

Describe the mode of action of a heterogeneous catalyst.

.....

.....

.....

..... [2]

[Total: 10]



- 4 (a) State what is meant by partition coefficient, K_{pc} .


.....

 [1]

- (b) Ethanedioic acid, $H_2C_2O_4$, is more soluble in water than in the organic solvent ethoxyethane.

2.70 g of ethanedioic acid is shaken with a mixture of 50 cm^3 of ethoxyethane and 50 cm^3 of water at 298 K and left until there is no further change in concentrations.

The ethoxyethane layer requires 16.5 cm^3 of 0.200 mol dm^{-3} NaOH(aq) for complete neutralisation.



Calculate the partition coefficient, K_{pc} , for ethanedioic acid between ethoxyethane and water.

$$K_{pc} = \dots\dots\dots [3]$$

- (c) The formula of ethanedioic acid is $HOOC-COOH$.

The formula of butanedioic acid is $HOOCCH_2CH_2COOH$.

Both acids are soluble in ethoxyethane and in water.

Suggest how the value of K_{pc} of butanedioic acid between ethoxyethane and water compares to the value of K_{pc} for ethanedioic acid between ethoxyethane and water.

Explain your answer.

.....

 [2]

[Total: 6]



- 5 (a) Define standard electrode potential, $E^\ominus$, including a description of standard conditions.

.....

.....

..... [2]

- (b) Table 5.1 shows the standard electrode potentials for three redox systems.

Table 5.1

redox system	electrode reaction	$E^\ominus/\text{V}$
1	$\text{Fe}^{3+} + \text{e}^- \rightleftharpoons \text{Fe}^{2+}$	+0.77
2	$\text{NO}_3^- + 10\text{H}^+ + 8\text{e}^- \rightleftharpoons \text{NH}_4^+ + 3\text{H}_2\text{O}$	+0.87
3	$\text{Zn}^{2+} + 2\text{e}^- \rightleftharpoons \text{Zn}$	-0.76

- (i) Use the information in Table 5.1 to write equations for **two** reactions that are feasible.

Explain why these reactions are feasible.

equation 1

equation 2

explanation

.....

..... [3]

- (ii) Suggest why the feasible reactions in 5(b)(i) may **not** take place under standard conditions.

.....

..... [1]



- (iii) An electrochemical cell is set up to measure $E^\ominus$ for redox system 2 in Table 5.1.

Complete Figure 5.1 to show a labelled diagram of this electrochemical cell.

Include all necessary substances and relevant pieces of apparatus needed to measure $E^\ominus$.

It is **not** necessary to state the conditions used.





Figure 5.1

[3]

- (c) A different electrochemical cell is set up. The cell reaction for this electrochemical cell is shown.



Calculate the standard cell potential, $E_{\text{cell}}^\ominus$, for this cell reaction.

$$E_{\text{cell}}^\ominus = \dots\dots\dots \text{ V [2]}$$

[Total: 11]

6 (a) Phenol reacts with $\text{Br}_2(\text{aq})$.

- (i) State **all** the expected observations when $\text{Br}_2(\text{aq})$ is added dropwise to a solution of phenol.

.....
..... [1]

- (ii) Draw the structure of the organic product formed when phenol reacts with an excess of $\text{Br}_2(\text{aq})$.

[1]

- (b) Explain the relative acidities of phenol and water.

.....
.....
.....
..... [2]



(c) Compound **C** can undergo both addition and condensation polymerisation.




Figure 6.1

(i) Draw **one** repeat unit of the addition polymer formed from **C**.

[1]

(ii) Draw **two** repeat units of the condensation polymer formed from **C**.

The new functional group formed should be displayed.

[2]

(d) **C** can react with LiAlH_4 . **C** can also react with an excess of $\text{H}_2(\text{g})$ with a nickel catalyst.

Draw the structure of the organic product formed in each reaction in Figure 6.2.




Figure 6.2

[2]



(e) Compound **F** can be synthesised from phenol in three steps, as shown in Figure 6.3.




Figure 6.3

(i) Draw the structures of compounds **D** and **E** in the boxes in Figure 6.3. [2]

(ii) Suggest reagents and conditions for steps 1 to 3 in Figure 6.3.

step 1

step 2

step 3

[3]

[Total: 14]



- 7 (a) Alanine, $\text{CH}_3\text{CH}(\text{NH}_2)\text{COOH}$, is an amino acid that contains a chiral centre.

Different synthetic routes to produce alanine can lead to the formation of an optically active sample or a racemic mixture.

- (i) Describe what is meant by optically active.

.....
..... [1]

- (ii) Describe what is meant by racemic mixture.

.....
..... [1]

- (b) Alanine has an isoelectric point of 6.11.

- (i) Explain what is meant by isoelectric point.

.....
..... [1]

- (ii) Draw the structure of alanine at pH 10.

[1]

- (c) Alanine molecules can react together to form the tripeptide Ala–Ala–Ala.

Draw the structure for this tripeptide. The functional groups formed should be displayed.

[2]



(d) Compound **J** can be synthesised from alanine in three steps, as shown in Figure 7.1.




Figure 7.1

- (i) Draw the structures of compounds **G** and **H** in the boxes in Figure 7.1. [2]
- (ii) Suggest reagents and conditions for steps 1 to 3 in Figure 7.1.

step 1

step 2

step 3 [3]

[Total: 11]

- 8 (a) Give the full name of the substance used in NMR spectroscopy as the standard for chemical shift measurements.

..... [1]

- (b) Compound **K** is analysed by carbon-13 NMR and proton (^1H) NMR spectroscopy in D_2O .




Figure 8.1

- (i) Predict the number of peaks in the carbon-13 NMR spectrum of **K**.

..... [1]

- (ii) The proton (^1H) NMR spectrum of **K** in D_2O shows four peaks, for the proton environments labelled **w**, **x**, **y** and **z** on the structure of **K** in Figure 8.2.




Figure 8.2

Complete Table 8.1 to identify the splitting patterns for proton environments **w**, **x**, **y** and **z**.

Table 8.1

proton environment	name of splitting pattern
w	
x	
y	
z	

[2]



- (iii) A separate sample of **K** is dissolved in CDCl_3 and analysed using proton (^1H) NMR spectroscopy.

The proton (^1H) NMR spectrum of **K** dissolved in CDCl_3 contains one more peak than the spectrum of **K** dissolved in D_2O .

Explain why there is one more peak, and suggest the chemical shift range in which it occurs.

.....

.....

.....

chemical shift range $\delta =$ ppm
[2]

Table 8.2

environment of proton	example	chemical shift range δ/ppm
alkane	$-\text{CH}_3$, $-\text{CH}_2-$, $>\text{CH}-$	0.9–1.7
alkyl next to $\text{C}=\text{O}$	$\text{CH}_3-\text{C}=\text{O}$, $-\text{CH}_2-\text{C}=\text{O}$, $>\text{CH}-\text{C}=\text{O}$	2.2–3.0
alkyl next to aromatic ring	CH_3-Ar , $-\text{CH}_2-\text{Ar}$, $>\text{CH}-\text{Ar}$	2.3–3.0
alkyl next to electronegative atom	CH_3-O , $-\text{CH}_2-\text{O}$, $-\text{CH}_2-\text{Cl}$	3.2–4.0
attached to alkene	$=\text{CHR}$	4.5–6.0
attached to aromatic ring	$\text{H}-\text{Ar}$	6.0–9.0
aldehyde	HCOR	9.3–10.5
alcohol	ROH	0.5–6.0
phenol	$\text{Ar}-\text{OH}$	4.5–7.0
carboxylic acid	RCOOH	9.0–13.0
alkyl amine	$\text{R}-\text{NH}-$	1.0–5.0
aryl amine	$\text{Ar}-\text{NH}_2$	3.0–6.0
amide	RCONHR	5.0–12.0

(c) Compound **K** reacts with an excess of hot NaOH(aq).






Figure 8.3

The products are isolated from the reaction mixture at pH 10.

Draw the structures of the **two** organic products of the complete alkaline hydrolysis of **K**.



[2]

[Total: 8]





Turn over





9 (a) Compound **P** and compound **Q** are isomers.




Figure 9.1

State the relative basicities of **P**, **Q** and ammonia.

Explain your answer.

..... > >
 most basic least basic

explanation

.....

.....

.....

.....

.....

.....

[4]



(b) **P** can be prepared from methylbenzene in two steps, as shown in Figure 9.2.




Figure 9.2

(i) Suggest reagents and conditions for steps 1 and 2 in Figure 9.2.

step 1

step 2 [2]

(ii) Complete the equation in Figure 9.3 for step 2.

Use [H] to represent an atom of hydrogen from the reducing agent.




Figure 9.3

[1]

(c) Figure 9.4 shows a two-step synthesis of azo dye **N**.




Figure 9.4

- (i) Draw the structure of compound **M** in the box in Figure 9.4. [1]
- (ii) Give the reagents and conditions for steps 1 and 2 in Figure 9.4.

step 1

step 2 [2]

[Total: 10]





Important values, constants and standards

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.022 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ (4.18 J g ⁻¹ K ⁻¹)

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (Cambridge University Press & Assessment) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in our Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge International Education is the name of our awarding body and a part of Cambridge University Press & Assessment, which is a department of the University of Cambridge.



The Periodic Table of Elements

Group																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																											
1	2	1 H hydrogen 1.0														13	14	15	16	17	18																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																						
		Key																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																									
		atomic number atomic symbol name relative atomic mass																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																									
3	Li lithium 6.9	4	Be beryllium 9.0															5	6	7	8	9	10	11	12	13	14	15	16	17	18																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																												
11	Na sodium 23.0	12	Mg magnesium 24.3															13	14	15	16	17	18																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																				
																		19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57–71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89–103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299	300	301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320	321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340	341	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360	361	362	363	364	365	366	367	368	369	370	371	372	373	374	375	376	377	378	379	380	381	382	383	384	385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400	401	402	403	404	405	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420	421	422	423	424	425	426	427	428	429	430	431	432	433	434	435	436	437	438	439	440	441	442	443	444	445	446	447	448	449	450	451	452	453	454	455	456	457	458	459	460	461	462	463	464	465	466	467	468	469	470	471	472	473	474	475	476	477	478	479	480	481	482	483	484	485	486	487	488	489	490	491	492	493	494	495	496	497	498	499	500	501	502	503	504	505	506	507	508	509	510	511	512	513	514	515	516	517	518	519	520	521	522	523	524	525	526	527	528	529	530	531	532	533	534	535	536	537	538	539	540	541	542	543	544	545	546	547	548	549	550	551	552	553	554	555	556	557	558	559	560	561	562	563	564	565	566	567	568	569	570	571	572	573	574	575	576	577	578	579	580	581	582	583	584	585	586	587	588	589	590	591	592	593	594	595	596	597	598	599	600	601	602	603	604	605	606	607	608	609	610	611	612	613	614	615	616	617	618	619	620	621	622	623	624	625	626	627	628	629	630	631	632	633	634	635	636	637	638	639	640	641	642	643	644	645	646	647	648	649	650	651	652	653	654	655	656	657	658	659	660	661	662	663	664	665	666	667	668	669	670	671	672	673	674	675	676	677	678	679	680	681	682	683	684	685	686	687	688	689	690	691	692	693	694	695	696	697	698	699	700	701	702	703	704	705	706	707	708	709	710	711	712	713	714	715	716	717	718	719	720	721	722	723	724	725	726	727	728	729	730	731	732	733	734	735	736	737	738	739	740	741	742	743	744	745	746	747	748	749	750	751	752	753	754	755	756	757	758	759	760	761	762	763	764	765	766	767	768	769	770	771	772	773	774	775	776	777	778	779	780	781	782	783	784	785	786	787	788	789	790	791	792	793	794	795	796	797	798	799	800	801	802	803	804	805	806	807	808	809	810	811	812	813	814	815	816	817	818	819	820	821	822	823	824	825	826	827	828	829	830	831	832	833	834	835	836	837	838	839	840	841	842	843	844	845	846	847	848	849	850	851	852	853	854	855	856	857	858	859	860	861	862	863	864	865	866	867	868	869	870	871	872	873	874	875	876	877	878	879	880	881	882	883	884	885	886	887	888	889	890	891	892	893	894	895	896	897	898	899	900	901	902	903	904	905	906	907	908	909	910	911	912	913	914	915	916	917	918	919	920	921	922	923	924	925	926	927	928	929	930	931	932	933	934	935	936	937	938	939	940	941	942	943	944	945	946	947	948	949	950	951	952	953	954	955	956	957	958	959	960	961	962	963	964	965	966	967	968	969	970	971	972	973	974	975	976	977	978	979	980	981	982	983	984	985	986	987	988	989	990	991	992	993	994	995	996	997	998	999	1000

lanthanoids

57La lanthanum 138.9	58Ce cerium 140.1	59Pr praseodymium 140.9	60Nd neodymium 144.2	61Pm promethium —	62Sm samarium 150.4	63Eu europium 152.0	64Gd gadolinium 157.3	65Tb terbium 158.9	66Dy dysprosium 162.5	67Ho holmium 164.9	68Er erbium 167.3	69Tm thulium 168.9	70Yb ytterbium 173.1	71Lu lutetium 175.0
89Ac actinium —	90Th thorium 232.0	91Pa protactinium 231.0	92U uranium 238.0	93Np neptunium —	94Pu plutonium —	95Am americium —	96Cm curium —	97Bk berkelium —	98Cf californium —	99Es einsteinium —	100Fm fermium —	101Md mendelevium —	102No nobelium —	103Lr lawrencium —

actinoids